



# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 1




## MIND MAP : LEARNING MADE SIMPLE

### CHAPTER - 2



# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 3



## MIND MAP : LEARNING MADE SIMPLE

### CHAPTER - 4

$E_a$  and proper orientation of the molecules determine the criteria for an effective collision.

$$\text{Rate} = P Z_{AV} e^{-E_a/RT}$$

P is Steric or Probability factor

Collision in which molecules collide with sufficient kinetic energy and proper orientation for breaking of bonds and formation of new bonds.

Number of collisions per unit volume of reaction mixture  
Rate =  $Z_{AV} e^{-E_a/RT}$

Number of reacting species taking part in an elementary reaction colliding to bring out a reaction.

Appearance of products or disappearance of reactants over a long time interval.  
 $\frac{d [P]}{dt}$  = slope  
 $r_{av} = \frac{-d [R]}{dt}$  = -slope

Rate of change in concentration of reactant/product at a particular time  
 $r_{inst} = \frac{-d[R]}{dt} = \frac{+d[P]}{dt}$

Rate of disappearance of R  
Decrease in concentration of R  
 $= \frac{-\Delta[R]}{\Delta t}$

Concentration : Higher the concentration of reactants, faster is the rate of reaction  
Temperature : Increases with increase in temperature, becomes almost double with  $10^{\circ}\text{C}$  rise.  
Presence of Catalyst : Increases with a catalyst.  
Surface Area : Greater is the surface area, faster is the rate of reaction.  
Activation Energy : Lower the activation energy, faster is the reaction.

Rate Law  
Fetters's Heterogeneous  
Arrhenius Equation

Change in concentration of reactants or products in unit time; Unit : mol L<sup>-1</sup>s<sup>-1</sup> or atm s<sup>-1</sup>

### Chemical Kinetics

Rate Constant: Rate of a reaction when concentration in unitry  
of each of the reactants is unity.

Integrated Rate Equation  
Half-life Of A Reaction  
Pseudo First Order Reaction  
Rate Of A Reaction  
Order Of Chemical Reaction  
Rate Of A Reaction  
Collision Frequency


Order	Integrated Rate Law	Straight Plot	Units (K)	Unit
0	$kt = [R]_0 - [R]$	$[R]$ vs $t$	$\text{mol L}^{-1} \text{s}^{-1}$	$\text{mol L}^{-1} \text{s}^{-1}$
1	$kt = \ln \left\{ \frac{[R]_0}{[R]} \right\}$	$\ln [R]$ vs $t$	$\text{s}^{-1}$	$\text{s}^{-1}$

Order	Half-life
0	$t_{1/2} = \frac{0.693}{k}$
1	$t_{1/2} = \frac{[\text{R}]_0}{2k}$
2	$t_{1/2} = \frac{[\text{R}]_0}{4k}$

- Are not truly of first order but under certain conditions behave as first order reaction
- Acid hydrolysis of ethyl acetate
- Inversion of sugar




# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 5




# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 6







# MIND MAP : LEARNING MADE SIMPLE

CHAPTER - 7




# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 8




# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 9




CHAPTER - 10



# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 11




(i) Simple/symmetrical : Alkyl or aryl attached to O, are same.  
 (ii) Mixed/Unsymmetrical : Two groups are different.


Common name : Alkyl group + OH IUPAC name substituting e of alkane with suffix of

Common name : Terms ortho, meta and para phenols are used. IUPAC name : Dihydroxy derivatives as 1,2-, 1,3- and 1,4-benzenediol


Common name : alkyl/aryl groups in alphabetical order replaced by ether. IUPAC name : In alkyl/aryl group 'e' followed by ether, oxy followed by parent hydrocarbon.



**3. From Grignard reagent**



1. From haloarenes  
 $\text{Cl-C}_6\text{H}_4-\text{NaOH} \xrightarrow[300\text{ atm}]{623\text{ K}} \text{O}^{\text{Na}}-\text{C}_6\text{H}_4-\text{HCl} \xrightarrow{\text{H}_2\text{O}} \text{C}_6\text{H}_5\text{OH}$
2. From benzene sulphonic acid  
 $\text{SO}_3\text{H} \xrightarrow{\text{Oleum}} \text{C}_6\text{H}_5-\text{NaOH} \xrightarrow{\text{H}^+} \text{C}_6\text{H}_5\text{OH}$
3. From diazonium salts  
 $\text{NH}_2-\text{C}_6\text{H}_4-\text{NaO}_2 \xrightarrow{\text{HCl}} \text{N}_2\text{+} + \text{C}_6\text{H}_5\text{OH}$
4. From Cumene  
 $\text{CH}_3-\text{C}_6\text{H}_5-\text{O}_2 \xrightarrow{\text{O}} \text{CH}_3-\text{C}_6\text{H}_5-\text{O}-\text{OH} \xrightarrow{\text{H}^+} \text{C}_6\text{H}_5\text{OH} + \text{CH}_3\text{COCH}_3$



# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 12


**ALDEHYDES AND KETONES:**

(i) Physical:  
Boiling points are higher than hydrocarbons and ethers of comparable molecular masses.

(ii) Chemical : Nucleophilic addition reactions :

Aldehydes are more reactive than ketones due to steric and electronic reasons.

$\text{HCN} + \text{OH}^- \xrightleftharpoons{\text{H}_2\text{O}} \text{CN}^- + \text{H}_2\text{O}$



$\text{R}-\overset{\delta^+}{\underset{\text{C}=\text{O}^+}{\text{C}}}=\text{O} + \text{CN}^- \xrightarrow{\text{H}_2\text{O}} \text{R}-\overset{\delta^+}{\underset{\text{C}=\text{O}^+}{\text{C}}}=\text{O}-\text{CN} = \text{C}(=\text{O})-\text{CN}$

$\text{R}-\overset{\delta^+}{\underset{\text{C}=\text{O}^+}{\text{C}}}=\text{O} + \text{HCl} \xrightarrow{\text{H}_2\text{O}} \text{R}-\overset{\delta^+}{\underset{\text{C}=\text{O}^+}{\text{C}}}=\text{O}-\text{H} + \text{Cl}^-$

$\text{R}-\overset{\delta^+}{\underset{\text{C}=\text{O}^+}{\text{C}}}=\text{O} + \text{H}_2\text{O} \xrightarrow{\text{H}_2\text{O}} \text{R}-\overset{\delta^+}{\underset{\text{C}=\text{O}^+}{\text{C}}}=\text{O}-\text{H}_2\text{O}$

Reduction : (a) To alcohols – aldehydes and ketones reduce to primary and secondary alcohols respectively by  $\text{NaBH}_4$  or  $\text{LiAlH}_4$ .

(b) To hydrocarbons –

$\text{C=O} \xrightarrow{\text{Zn-Hg}/\text{HCl}} \text{CH}_2 + \text{H}_2\text{O}$  (Clemmensen Reduction)

$\text{C=O} \xrightarrow{\text{NH}_2\text{NH}/\text{H}_2\text{O}} \text{C=NNH}_2 \xrightarrow{\text{KOH/Ethylene glycol}/\text{Heat}} \text{CH}_2 + \text{N}_2$  (Wolf-Kishner)

Oxidation:  $\text{RCHO} \xrightarrow{\text{[O]}} \text{R-COOH}$

Fehling's test :  $\text{RCHO} + 2[\text{Ag}(\text{NH}_3)_2]^+ + 3\text{OH}^- \rightarrow \text{RCOO}^- + 2\text{Ag} + 2\text{H}_2\text{O} + 4\text{NH}_3$

$\text{RCHO} + 2\text{Cu}^{2+} + 5\text{OH}^- \rightarrow \text{RCOO}^- + \text{Cu}_2\text{O} + 3\text{H}_2\text{O}$

Haloform reaction:  $\text{R}-\overset{\delta^+}{\underset{\text{C}=\text{O}^+}{\text{C}}}=\text{O} \xrightarrow{\text{NaOCl}/\text{H}_2\text{O}} \text{R}-\text{C}(=\text{O})-\text{ONa} + \text{CH}_3\text{O}$

Reactions due to  $\alpha$ -hydrogen:

$2\text{CH}_3\text{CHO} \xrightarrow{\text{dil NaOH}/\Delta} \text{CH}_3-\overset{\text{CH}_3}{\underset{\text{OH}}{\text{C}}}=\text{CH}-\text{CHO} \xrightarrow{\text{H}_2\text{O}} \text{CH}_3-\text{CH}=\text{CH}-\text{CHO}$

$2\text{CH}_3\text{COCH}_3 \xrightarrow{\text{Ba(OH)}_2/\Delta} \text{CH}_3-\overset{\text{CH}_3}{\underset{\text{OH}}{\text{C}}}=\text{CH}_2-\text{COCH}_3 \xrightarrow{\text{H}_2\text{O}} \text{CH}_3-\overset{\text{CH}_3}{\underset{\text{OH}}{\text{C}}}=\text{CH}_2-\text{CO}-\text{CH}_3$

$\text{CH}_3\text{CHO} \xrightarrow{\text{NaOH}/\Delta} \text{CH}_3-\text{CH}=\text{CH}-\text{CHO} + \text{CH}_3-\text{CH}_2-\text{CH}=\text{C}-\text{CHO}$

Cannizzaro reaction :  $2\text{RCHO} + \text{conc KOH} \xrightarrow{\Delta} \text{CH}_3\text{OH} + \text{HCOOK}$

Electrophilic substitution reaction:

$\text{C}_6\text{H}_5\text{CHO} \xrightarrow{273-283\text{K}} \text{C}_6\text{H}_5\text{CO}$

Carboxylic acids:

(i) Physical: Higher boiling points than aldehydes, ketones or alcohols.  
Solubility decreases with increasing number of C atoms

(ii) Chemical :  $2\text{RCOOH} + 2\text{Na} \longrightarrow 2\text{RCOONa} + \text{H}_2$

Forms corresponding anhydride on heating with mineral acids

$\text{RCOOH} + \text{KOH} \xrightleftharpoons{\text{H}^+} \text{RCOOR} + \text{H}_2\text{O}$

$\text{RCOOH} + \text{PCl}_5 \xrightarrow{\text{H}_2\text{O}} \text{RCOCl} + \text{POCl}_3 + \text{HCl}$



$\text{CH}_3\text{COOH} + \text{NH}_3 \rightleftharpoons \text{CH}_3\text{COONH}_4 \xrightarrow{\Delta} \text{CH}_3\text{CONH}_2$

$\text{RCOOH} \xrightarrow{\text{B}_2\text{H}_6} \text{RCH}_2\text{OH}$

$\text{RCOONa} \xrightarrow{\text{NaOH} \& \text{CaO}/\text{Heat}} \text{R}-\text{H} + \text{Na}_2\text{CO}_3$


$\text{RCH}_2\text{COOH} \xrightarrow[\text{H}_2\text{O}]{\text{X/Red P}} \text{R}-\overset{\text{X}}{\underset{\text{C}}{\text{CH}}}=\text{COOH}$  (HVZ reaction)

$\text{CHO} \xrightarrow{\text{Conc HNO}_3} \text{CHO}_2$



# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 13





Preparation :  $C_6H_5NH_2 + NaNO_2 + 2HCl \xrightarrow{273-278K} C_6H_5N_2Cl + NaCl + 2H_2O$

Physical properties : Colourless crystalline solid, soluble in water; stable in cold but reacts with water on warming.



Chemical properties :

- Sandmeyer reaction:  $ArN_2^+ \xrightarrow{\substack{Cu_2Cl_2/HCl \\ Cu_2Br_2/HBr \\ CuCN/KCN}} ArBr + N_2$
- Gattermann reaction :  $ArN_2^+ \xrightarrow{\substack{Cu/HCl \\ Cu/HBr \\ ArBr + N_2 + CuX}}$
- Coupling reaction :  $ArN_2^+ + H_2O \xrightarrow{\substack{H^+ \\ H_3PO_2}} ArOH + N_2 + HCl$




# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 14




MIND MAP : LEARNING MADE SIMPLE CHAPTER - 15



# MIND MAP : LEARNING MADE SIMPLE

## CHAPTER - 16



**Purpose:**

- For their preservation.
- Enhancing their appeal.
- Adding nutritive value.

(a) Artificial Sweetening Agents : Natural sweeteners (sucrose), artificial sweeteners (Aspartame, Saccharin)

(b) Food Preservatives : Prevent spoilage of food due to microbial growth. (Table salt, sugar)

